

358

a)

x	$f(x) = 2x - 5$	(x, y)
0	$f(0) = 2 \cdot 0 - 5 = -5$	(0, -5)
1	$f(1) = 2 \cdot 1 - 5 = -3$	(1, -3)
2	$f(2) = 2 \cdot 2 - 5 = -1$	(2, -1)

b)

x	$g(x) = -2x - 2$	(x, y)
0	$g(0) = -2 \cdot 0 - 2 = -2$	(0, -2)
1	$g(1) = -2 \cdot 1 - 2 = -4$	(1, -4)
2	$g(2) = -2 \cdot 2 - 2 = -6$	(2, -6)


$$-2 - 2 = -4$$

c)

x	$h(x) = 3x + 2$	(x, y)
0	$h(0) = 3 \cdot 0 + 2 = 2$	(0, 2)
1	$h(1) = 3 \cdot 1 + 2 = 5$	(1, 5)
2	$h(2) = 3 \cdot 2 + 2 = 8$	(2, 8)
-1	$h(-1) = 3 \cdot (-1) + 2 = -1$	(-1, -1)

d)

x	$k(x) = -x - 3$	(x, y)
0	$k(0) = -0 - 3 = -3$	(0, -3)
1	$k(1) = -1 - 3 = -4$	(1, -4)
2	$k(2) = -2 - 3 = -5$	(2, -5)
-1	$k(-1) = -(-1) - 3 = 1 - 3 = -2$	(-1, -2)



359

360

5.93